

# Step-up to High Dose Fluticasone Furoate in Combination With Long-Acting Bronchodilator in Inadequately Controlled Asthma: The CAPTAIN Study

Poster No. 072

Hanania NA<sup>1</sup>, Kerwin E<sup>2</sup>, Pavord ID<sup>3</sup>, Kerstjens H<sup>4</sup>, Pascoe S<sup>5\*</sup>, Peachey G<sup>6</sup>, Fowler A<sup>6</sup>, Bailes Z<sup>6</sup>, Edwards D<sup>6</sup>, Sule N<sup>5</sup>, Barnes N<sup>7,8</sup>, Gardiner F<sup>6</sup>, Lee LA<sup>5</sup>

<sup>1</sup>Baylor College of Medicine, Houston, TX, USA; <sup>2</sup>Crisor LLC Research, Clinical Research Institute of Southern Oregon, Medford, OR, USA; <sup>3</sup>University of Oxford, Oxford, UK; <sup>4</sup>University of Groningen and University Medical Center Groningen, Groningen, the Netherlands; <sup>5</sup>GSK, Upper Providence, PA, USA; <sup>6</sup>GSK, Stockley Park West, Uxbridge, Middlesex, UK; <sup>7</sup>GSK, Brentford, Middlesex, UK; <sup>8</sup>Barts and the London School of Medicine and Dentistry, London, UK  
\*Affiliation at time of study


## Background

- Up to 50% of patients with asthma are inadequately controlled despite receiving dual therapy with an inhaled corticosteroid (ICS) and long-acting  $\beta_2$ -agonist (LABA).<sup>1-5</sup> Current guidelines for these patients recommend increasing the ICS dose in a stepwise manner.<sup>6</sup>
- Adding vilanterol (VI) to fluticasone furoate (FF) 100 mcg improves lung function and clinical outcomes in patients uncontrolled on ICS alone<sup>1</sup>; however, little evidence exists on the effect of doubling the ICS component in dual therapy.

## Objective

- This analysis of the CAPTAIN study reports on the effects of doubling the FF dose in FF/VI dual therapy in patients with inadequately controlled asthma on ICS/LABA therapy.

Figure 1. Study design



\*Provided BID as a fixed dose via DISKUS dry powder inhaler; †provided QD as a fixed-dose via the Ellipta dry powder inhaler; ‡all patients in the study had a safety follow-up contact approximately 7 days after the End of Study Visit (dependent on actual treatment date) or Early Withdrawal Visit; §responders were defined as patients with ≥0.5- or ≥4-point improvement from baseline in ACQ-7 and SGRQ total score, respectively; †safety endpoints reported here include incidence and type of adverse events (AEs), serious AEs (SAEs), and AEs of special interest (AESIs) relevant to ICS class effects. ACQ-6, Asthma Control Questionnaire-6; BID, twice daily; CFB, change from baseline; FF, fluticasone propionate; ITT, intent-to-treat; QD, once daily; R, randomization; SAL, salmeterol; SGRQ, St George's Respiratory Questionnaire

## References

- Bateman ED et al. *Thorax* 2014;69:312–9; 2. Bernstein DI, et al. *J Asthma* 2015;52:1073–83; 3. Davis J, et al. *J Asthma* 2019;56:332–40; 4. Lee LK, et al. *J Asthma* 2018;55:208–19; 5. Sulaiman S et al. *Eur Resp J* 2018;51:1701126; 6. Global Initiative for Asthma (GINA). Global strategy for asthma management and prevention. Available at: <https://ginasthma.org/wp-content/uploads/2019/06/GINA-2019-main-report-June-2019-wms.pdf> [last accessed January 2020].

## Disclosures

- This study was funded by GlaxoSmithKline (GSK study 205715, NCT02924688). ELLIPTA and DISKUS are owned by or licensed to the GSK group of companies.
- ZB, NB, DE, AF, LAL, GP, NS, and FG are employees of GSK hold stocks or shares in GSK. SP was an employee of GSK at the time of study and owns stocks in GSK. NAH reports receiving personal fees from AstraZeneca, Boehringer Ingelheim, Genentech, Genzyme, GSK, Mylan, Novartis, Regeneron, Sanofi, Sunovion, and for serving as an advisor or consultant. He also received research support from Astra Zeneca,

Boehringer Ingelheim and GSK. HK has received research grants and served on advisory boards for Boehringer Ingelheim, GSK, and Novartis. EK is an employee of Crisor LLC Research and has served on advisory boards, speaker panels or received travel reimbursement from Amphastar, AstraZeneca, Boehringer Ingelheim, Forest, GSK, Mylan, Novartis, Pearl, Sunovion, Teva and Theravance. EK has also conducted multicenter clinical research trials for approximately 40 pharmaceutical companies. IDP has received speaker's fees, payments for organizing education events, honoraria for attending advisory panels, sponsorship to attend international scientific meetings, research grants or payments to support FDA approval meetings from Aerocrine, Almirall, AstraZeneca, Boehringer Ingelheim, Chiesi, Circassia, Genentech, GSK, Knopp, Merck, Novartis, Sanofi/Regeneron and Teva.

IDP acted as an expert witness for a patent dispute involving AstraZeneca and Teva, is a Co-patent holder for the Leicester Cough Questionnaire, and has received payments for use of the Leicester Cough Questionnaire in clinical trials from Bayer, Insmad, and Merck.

- Editorial support (in the form of writing assistance, including development of the initial draft based on author direction, assembling tables and figures, collating authors' comments, grammatical editing, and referencing) was provided by Philip Chapman, of Fishawack Indicia Ltd, UK, and was funded by GSK.

Please find the online version of this poster by scanning the QR code or via [http://tago.ca/AAAA1\\_8](http://tago.ca/AAAA1_8)



## Results


- 2436 patients were enrolled in the ITT population. Clinically meaningful improvements from baseline in mean (standard deviation) trough FEV<sub>1</sub> (287 [356] mL) and ACQ-7 score (-0.674 [0.705]) were seen during the 3-week run-in period with FSC and 2-week stabilization period with FF/VI; improvements with FF/VI (162 [290] mL; -0.387 [0.575]) were greater than FSC (126 [307] mL; -0.283 [0.524]).
- Baseline demographics and clinical characteristics were similar between FF/VI treatment groups (Table 1).
- At Week 24, mean change from baseline in trough FEV<sub>1</sub> improved with FF/VI 200/25 mcg compared with FF/VI 100/25 mcg (Figure 2); improvements were also observed in clinic FEV<sub>1</sub> at 3 hours post-treatment (treatment difference: 36 mL [95% CI: -8, 81]).
- In total, 688 patients (28%) experienced moderate/severe exacerbations and 73 and 106 severe exacerbations were reported for FF/VI 200/25 mcg and FF/VI 100/25 mcg, respectively, over 52 weeks. FF/VI 200/25 mcg also resulted in a lower risk of moderate/severe exacerbation (adjusted rate ratio [95% CI]: 0.65 [0.50, 0.85]) versus FF/VI 100/25 mcg (annualized rate FF/VI 100/25 mcg 0.87, FF/VI 200/25 mcg 0.57). Time to first moderate/severe exacerbation is shown in Figure 3.
- Greater improvements in ACQ-7 total score and more ACQ-7 responders at Week 24 (Figure 4) were seen with FF/VI 200/25 mcg compared with FF/VI 100/25 mcg. No difference in SGRQ total score and proportion of responders was observed between both treatments.
- The safety profile for both treatments was similar (Table 2).

Table 1. Baseline demographics and clinical characteristics (ITT population)

	FF/VI 100/25 mcg (N=407)	FF/VI 200/25 mcg (N=406)
Age, years, mean (SD)	53.3 (13.03)	53.9 (13.30)
Male, n (%)	153 (38)	154 (38)
Onset age of asthma, years, mean (SD)	32.8 (19.09)	33.2 (18.93)
BMI (kg/m <sup>2</sup> ), mean (SD)	29.29 (6.08)	29.38 (6.29)
Total number of exacerbations in 12 months prior to screening, n (%)		
0	62 (15)	62 (15)
1	219 (54)	251 (62)
≥2	126 (31)	93 (23)
Former smokers, n (%)	69 (17)	69 (17)
Eosinophils at screening (10 <sup>9</sup> /L), mean (SD)*	0.327 (0.274)	0.305 (0.239)
Pre-bronchodilator FEV <sub>1</sub> , % predicted, mean (SD) (screening) <sup>†</sup>	58.24 (13.06)	58.66 (13.20)
ACQ-6 score, mean (SD) <sup>‡</sup> (screening)	2.48 (0.61)	2.53 (0.66)


\*FF/VI 100/25 mcg, n=382 and FF/VI 200/25 mcg, n=389; †FF/VI 100/25 mcg, n=402 and FF/VI 200/25 mcg, n=401; ‡FF/VI 100/25 mcg, n=406. BMI, body mass index; SD, standard deviation

Figure 2. LS mean change from baseline in clinic trough FEV<sub>1</sub> at Week 24 (ITT population)




Analysis performed using mixed model repeated measures with covariates of treatment, age, sex, region, baseline value, pre-study ICS dosage at screening, and visit, interaction terms for baseline value by visit and treatment by visit. LS, least squares

Figure 3. Time to first moderate/severe exacerbation (ITT population)



Analysis performed using a Cox proportional hazards model with covariates of treatment, age, sex, region, pre-study ICS dosage at screening, severe asthma exacerbations in the previous year (0, 1, ≥2). HR, hazard ratio

Figure 4. (A) LS mean change from baseline and (B) responder rates for ACQ-7 total score at Week 24 (ITT population)



Responder threshold is defined as an improvement (decrease) of ≥0.5 points from baseline. Analysis performed using mixed model repeated measures (for A) and generalized linear mixed model with a logit link function (for B) with covariates of treatment, age, sex, region, baseline value, pre-study ICS dosage at screening, and visit, interaction terms for baseline value by visit and treatment by visit.

## Conclusions

- Clinically significant improvements from screening in trough FEV<sub>1</sub> and ACQ score were observed during the run-in and stabilization period with FSC and FF/VI 100/25 mcg.
- Doubling the FF dose provided modest lung function improvements and a 35% reduction in annualized moderate/severe exacerbation rates. Trends for improvements in asthma control were noted and there was no increase in the risk of AEs.
- These results demonstrate the dose–response effect of FF/VI on clinical outcomes and suggest that increasing the FF dose may be a suitable treatment-escalation option for patients uncontrolled with ICS/LABA therapy.

Table 2. On-treatment AEs and AESIs relevant to ICS class effects (ITT population)

n (%)	FF/VI 100/25 mcg (N=407)	FF/VI 200/25 mcg (N=406)
Any AE	258 (63)	210 (52)
AE leading to discontinuation	11 (3)	5 (1)
Drug-related AE	21 (5)	17 (4)
SAE	25 (6)	21 (5)
<b>AESIs relevant to ICS class effects</b>		
Decreased BMD and associated fractures	5 (1)	2 (<1)
Hyperglycemia/new-onset DM	12 (3)	8 (2)
Infective pneumonia	7 (2)	7 (2)
LRTI excluding infective pneumonia	20 (5)	25 (6)
Local steroid effects	12 (3)	17 (4)

Note: Preferred terms may have contributed to more than one special interest group. AESIs were counted in each special interest group in which they appeared. BMD, bone mineral density; DM, diabetes mellitus; LRTI, lower respiratory tract infection